

Further, our findings indicate that the role of TN-GnRH3 may be closely associated with female preference for a visually recognized individual. Individual recognition is central to many types of cooperative interactions (23), social decision-making in pair-bonding (4), kin recognition (1, 24), and social hierarchy (25). The present findings shed new light on the importance of TN-GnRH3 neurons for female preference, as well as for individual recognition, for social neuroscience.

#### References and Notes

1. C. Brown, K. N. Laland, J. Krause, *Fish Cognition and Behaviour* (Blackwell, Oxford, 2006).
2. M. N. Verzijden et al., *Trends Ecol. Evol.* **27**, 511–519 (2012).
3. K. A. Hughes, L. Du, F. H. Rodd, D. N. Reznick, *Anim. Behav.* **58**, 907–916 (1999).
4. L. J. Young, Z. Wang, *Nat. Neurosci.* **7**, 1048–1054 (2004).
5. M. Kasahara et al., *Nature* **447**, 714–719 (2007).
6. T. Ono, T. Uematsu, *J. Fac. Sci. Hokkaido Univ.* **13**, 197 (1957).
7. R. O. Walter, J. B. Hamilton, *Anim. Behav.* **18**, 125–127 (1970).
8. T. Sasado et al., *Dev. Biol.* **320**, 328–339 (2008).
9. Y. Toba, J. D. Tiong, Q. Ma, S. Wray, *Dev. Neurobiol.* **68**, 487–503 (2008).
10. O. Palevitch et al., *Gen. Comp. Endocrinol.* **165**, 262–268 (2010).
11. K. Okubo et al., *Endocrinology* **147**, 1076–1084 (2006).
12. H. Abe, Y. Oka, *J. Reprod. Dev.* **57**, 665–674 (2011).
13. H. Abe, Y. Oka, *J. Neurophysiol.* **83**, 3196–3200 (2000).
14. A. L. Kyle, B. G. Luo, T. H. Magnus, W. K. Stell, *Cell Tissue Res.* **280**, 605–615 (1995).
15. Y. Akazome, S. Kanda, Y. Oka, *J. Neuroendocrinol.* **23**, 570–576 (2011).
16. K. W. Sockman, G. F. Ball, *Dev. Neurobiol.* **69**, 339–349 (2009).
17. W. Wilczynski, M. J. Ryan, *Curr. Opin. Neurobiol.* **20**, 754–763 (2010).

**Acknowledgments:** We thank the National BioResource Project Medaka for supplying the medaka strains; Y. Wakamatsu for the STII strains; Solution Oriented Research for Science and Technology (SORST) Kondoh Research Team for *cxcr4*(kazura) and *cxcr7*(yanagi) mutant strains; Y. Nagahama for Tg(*gnrh1:gfp*), Tg(*gnrh2:gfp*), and Tg(*gnrh3:gfp*); T. Sasado,

S. Nonaka, H. Otsuna, S. Kanda, T. Yamamoto, E. Yamamoto, T. Karigo, and I. Hara, for technical assistance; T. Hiraki for NPFF immunostaining; and S. Nakagawa, F. Loosti, S. Furutani-Seiki, H. Kondoh, and F. Gabbiani for discussion. This work was supported by National Institute for Basic Biology Priority Collaborative Research Project (10-104), Grant-in-Aid for Scientific Research (B) 23300115, Grant-in-Aid for a Japan Society for Promotion of Science (JSPS) postdoctoral SPD fellowship (to T.O.), the Cell Science Research Foundation (to T.O.), Comprehensive Brain Science Network (to T.O.), and Grant-in-Aid for JSPS fellows (to S. Yokoi, Y.S., H.I. and Y.I.). There are MTAs associated with *cxcr4*(kazura), *cxcr7*(yanagi), d-r-Tg (*olvas:GFP*) medaka and ST-III medaka as provided by the National BioResource Project (NBRP).

#### Supplementary Materials

www.sciencemag.org/content/343/6166/91/suppl/DC1  
Materials and Methods  
Figs. S1 to S19  
Tables S1 and S2  
Movies S1 to S3  
References (18–39)

15 August 2013; accepted 18 November 2013  
10.1126/science.1244724

## Pregnenolone Can Protect the Brain from Cannabis Intoxication

Monique Vallée,<sup>1,2†</sup> Sergio Vitiello,<sup>1,2\*</sup> Luigi Bellocchio,<sup>1,2\*</sup> Etienne Hébert-Chatelain,<sup>1,2\*</sup> Stéphanie Monlezun,<sup>3\*</sup> Elena Martin-Garcia,<sup>4</sup> Fernando Kasantz,<sup>1,2</sup> Gemma L. Baillie,<sup>5,7</sup> Francesca Panin,<sup>1,2</sup> Adeline Cathala,<sup>1,2</sup> Valérie Roullot-Lacarrière,<sup>1,2</sup> Sandy Fabre,<sup>3</sup> Dow P. Hurst,<sup>6</sup> Diane L. Lynch,<sup>6</sup> Derek M. Shore,<sup>6</sup> Véronique Deroche-Gamonet,<sup>1,2</sup> Umberto Spampinato,<sup>1,2</sup> Jean-Michel Revest,<sup>1,2</sup> Rafael Maldonado,<sup>4</sup> Patricia H. Reggio,<sup>6</sup> Ruth A. Ross,<sup>5,7</sup> Giovanni Marsicano,<sup>1,2</sup> Pier Vincenzo Piazza<sup>1,2†‡</sup>

Pregnenolone is considered the inactive precursor of all steroid hormones, and its potential functional effects have been largely uninvestigated. The administration of the main active principle of *Cannabis sativa* (marijuana),  $\Delta^9$ -tetrahydrocannabinol (THC), substantially increases the synthesis of pregnenolone in the brain via activation of the type-1 cannabinoid (CB<sub>1</sub>) receptor. Pregnenolone then, acting as a signaling-specific inhibitor of the CB<sub>1</sub> receptor, reduces several effects of THC. This negative feedback mediated by pregnenolone reveals a previously unknown paracrine/autocrine loop protecting the brain from CB<sub>1</sub> receptor overactivation that could open an unforeseen approach for the treatment of cannabis intoxication and addiction.

**S**teroid hormones are important modulators of brain activity and behavior (1–4). Steroids play crucial roles in regulating physiological activities such as food intake, wakening, reproduc-

tion, and sexual behavior and participate in the regulation of mood and memory. Steroids also facilitate coping with stress and have been implicated in stress-related pathologies (1–4).

Although the most-studied steroids are produced in the periphery, some of them, named neurosteroids, are also synthesized directly in the brain (5, 6) from the putatively inactive precursor pregnenolone (3 $\beta$ -hydroxypregn-5-en-20-one) (5). Active neurosteroids, such as pregnenolone sulfate (20-oxo-5-pregnen-3 $\beta$ -yl sulfate), allopregnanolone (3 $\alpha$ -hydroxy-5 $\alpha$ -pregnan-20-one), and DHEA (3 $\beta$ -hydroxyandrost-5-en-17-one), have been implicated in the regulation of mood and cognitive activities, and their decline has been associated with aging-related impairments (5, 7).

We investigated the involvement of neurosteroids in addiction by studying the effects of the major classes of drugs of abuse on their production in the brains of rats and mice. Concentrations of brain steroids were analyzed using gas chromatog-

raphy coupled to mass spectrometry (8, 9), which allows measuring, in the same sample, pregnenolone, DHEA, testosterone (17 $\beta$ -hydroxyandrost-4-en-3-one) and its metabolite DHT (dihydrotestosterone; 17 $\beta$ -hydroxy-5 $\alpha$ -androst-3-one), and the three stereoisomers pregnanolone (3 $\alpha$ -hydroxy-5 $\beta$ -pregnan-20-one), allopregnanolone, and epiallopregnanolone (3 $\beta$ -hydroxy-5 $\alpha$ -pregnan-20-one). As shown for the ventral striatum (the nucleus accumbens, NAc), in the brain of Wistar rats, basal levels were approximately 1 ng per gram of tissue (ng/g) for pregnenolone and testosterone, around 0.4 ng/g for allopregnanolone and DHT, and only traces of epiallopregnanolone (<0.2 ng/g) were found (Fig. 1A). In C57BL/6N mice, the highest concentrations were found for pregnenolone and epiallopregnanolone, and the lowest concentrations were observed for testosterone. DHT was undetectable (fig. S1A). In both rat and mice brains, DHEA and pregnanolone were undetectable under basal conditions and after the administration of drugs.

Representative compounds of the major classes of drugs of abuse were injected into Wistar rats at doses corresponding approximately to the median effective dose (ED<sub>50</sub>) for most of their unconditional behavioral effects: cocaine [20 mg per kilogram of body weight (mg/kg)], morphine (2 mg/kg), nicotine (0.4 mg/kg), alcohol (1 g/kg), and the main active principle of marijuana (*Cannabis sativa*),  $\Delta^9$ -tetrahydrocannabinol (THC) (3 mg/kg) (10). The increase in pregnenolone (Fig. 1B and table S1) induced by THC (around 1500%) was several times higher and longer-lasting (2 hours) than the one induced by the other drugs (around 300% and 30 min). Dose-response studies showed a maximal THC-induced increase in pregnenolone of approximately 3000% in both Wistar rats (Fig. 1C) and C57BL/6N mice (fig. S1).

The effects of THC on pregnenolone-derived neurosteroids were not statistically significant in mice (fig. S1). In rats (Fig. 1C), a statistically sig-


<sup>1</sup>INSERM, Neurocentre Magendie, Physiopathologie de la Plasticité Neuronale, U862, F-33000 Bordeaux, France. <sup>2</sup>Université de Bordeaux, Neurocentre Magendie, Physiopathologie de la Plasticité Neuronale, U862, F-33000 Bordeaux, France. <sup>3</sup>Alienor Farma Parc Scientifique Unitec1, 2 Allée du Doyen Georges Brus, 33600 Pessac, France. <sup>4</sup>Laboratory of Neuropharmacology, Department of Experimental and Health Sciences, Universitat Pompeu Fabra, Parc de Recerca Biomèdica de Barcelona, Calle Dr. Aiguader 88, Barcelona 08003, Spain. <sup>5</sup>Kosterlitz Centre for Therapeutics, Institute of Medical Sciences, University of Aberdeen, Aberdeen AB25 2ZD, UK. <sup>6</sup>Department of Chemistry and Biochemistry, University of North Carolina at Greensboro, Greensboro, NC, USA. <sup>7</sup>Present address: Department of Pharmacology and Toxicology, University of Toronto, Medical Sciences Building, 1 King's College Circle, Toronto, Ontario, Canada M5S 1A8.

\*These authors contributed equally to this work.


†These authors equally supervised this work.

‡Corresponding author. E-mail: pier-vincenzo.piazza@inserm.fr

**Fig. 1. THC increases pregnenolone levels by activating the CB<sub>1</sub> receptor.** (A) Basal levels of pregnenolone (PREG), allopregnanolone (ALLO), epiallopregnanolone (EPI), testosterone (T), and dihydrotestosterone (DHT) in the NAC. (B) Compared to the major classes of drugs of abuse, cocaine [20 mg/kg, administered intraperitoneally (ip)], morphine (2 mg/kg, ip), nicotine (0.4 mg/kg, ip), and ethanol (1 g/kg, ip), THC (3 mg/kg, ip) induced the highest increase in pregnenolone concentrations in the NAC. The arrow indicates the time of drug injection. (C) THC dose-dependently increased [F(6,30) = 17.2,  $P < 0.001$ ] pregnenolone concentrations in the NAC, with minor effects on pregnenolone-derived downstream steroids. (D and E) THC at 9 mg/kg differently increased pregnenolone concentrations in brain structures and peripheral tissues: the prefrontal cortex (FCX), NAC, dorsal striatum (STR), hippocampus (HPC), thalamus (THA), hypothalamus (HYP), ventral midbrain (VMB), sensory motor cortex (CX), cerebellum (CB), spinal cord (SPI), kidney (KID), liver (LIV), spleen (SPL), lung (LUN), intestine (INT), muscle (MUS), white adipose tissue (WAT), testis (TES), and plasma. (F) In the NAC, the ip injection of the CB<sub>1</sub> agonists HU210 and WIN 55,212-2 dose-dependently increased pregnenolone levels [analysis of variance (ANOVA),  $P < 0.001$  in all cases]. The CB<sub>2</sub> agonist JWH-133 had non-statistically significant effects. (G) The increase in pregnenolone concentrations induced by THC (3 mg/kg, ip) in the NAC was abolished by the CB<sub>1</sub> antagonist AM251 (8 mg/kg, ip) injected 30 min before THC. THC (12 mg/kg, ip) induced an increase in pregnenolone levels in the NAC of wild-type mice but not in KO mice with a (H) complete (CB<sub>1</sub><sup>−/−</sup>) or (I) neuron-specific (D<sub>1</sub>-CB<sub>1</sub><sup>−/−</sup>)



**Fig. 2. THC can increase pregnenolone synthesis through proteins involved in neurosteroidogenesis.** Schematic representation of (A) the proposed molecular mechanism and (B) the protocol used. (C) Representative Western blots and (D) densitometric quantification of NAC expression of cytochrome P450scc, StAR, P-HSL<sup>Ser660</sup>, HSL, and  $\beta$ III-tubulin proteins, in Wistar rats ip injected with THC (9 mg/kg) after treatment with SL327 or vehicle. 15 min after THC administration, the levels of cytochrome P450scc increased via an Erk1/2<sup>MAPK</sup>-dependent mechanism. 30 min after THC administration, with an Erk1/2<sup>MAPK</sup>-independent mechanism, cytochrome P450scc was still increased and HSL was activated by phosphorylation. THC administration did not modify the levels of StAR proteins. Data are expressed as mean  $\pm$  SEM ( $n = 5$  to 7 animals per group). OD, optical density. \* $P < 0.05$ , \*\*\* $P < 0.001$  in comparison to vehicle-treated rats (white and white-striped bars). # $P < 0.05$ , ### $P < 0.005$ , ### $P < 0.001$  in comparison to THC-treated rats (black and black-striped bars). Fisher's protected least significant difference post hoc test was used after ANOVA.



deletion of the CB<sub>1</sub> receptor. Data are expressed as mean  $\pm$  SEM ( $n = 6$  to 12 animals per group). \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$  compared to animals that did not receive THC. All experiments except (H) and (I) were performed in Wistar rats.

nificant effect was observed only for allopregnanolone and epiallopregnanolone. However, even at the highest dose of THC (9 mg/kg), the increase in these pregnenolone metabolites (Fig. 1C) was several times lower than the increase observed in pregnenolone (in ng/g: allopregnanolone =  $0.73 \pm 0.14$ , epiallopregnanolone =  $0.40 \pm 0.08$ , and pregnenolone =  $28.38 \pm 3.16$ ).

The highest THC-induced increase in pregnenolone in Wistar rats (Fig. 1D) was observed in the NAc, prefrontal cortex, striatum, and thalamus and the lowest in the spinal cord, ventral midbrain, sensory motor cortex, and peripheral tissues such as the kidney, spleen, lung, and white fat (Fig. 1E). In the liver, gastrointestinal tract, muscle, testis, and plasma, THC had no significant effect.

The effects of THC in the brain are mainly mediated by the type-1 cannabinoid (CB<sub>1</sub>) receptor (11–13). In the NAc, similarly to THC, injection of Wistar rats with the CB<sub>1</sub>/CB<sub>2</sub> cannabinoid receptor agonists HU210 or WIN55,212-2 increased pregnenolone levels (Fig. 1F) at doses that are in line with their respective affinities for the CB<sub>1</sub> receptor (14). The CB<sub>2</sub>-selective agonist JWH133 had no significant effect (Fig. 1F). THC effects on pregnenolone in the NAc were suppressed (i) by the CB<sub>1</sub>-selective antagonist AM251 in Wistar rats (Fig. 1G); (ii) in constitutive CB<sub>1</sub> receptor knockout (KO) mice (CB<sub>1</sub><sup>−/−</sup>, Fig. 1H), which lack CB<sub>1</sub> receptors in all cell types; and (iii) in conditional mutant mice (15), which lack CB<sub>1</sub> receptors in the majority of striatal  $\gamma$ -aminobutyric acid (GABAergic) spiny neurons [the ones containing the dopamine receptor D<sub>1</sub> (D<sub>1</sub>-CB<sub>1</sub><sup>−/−</sup>)] (Fig. 1I) (9). The CB<sub>1</sub>-KO mice were from mice strains in which the C57BL/6N genotype was predominant (six or seven backcrossing generations). These data indicate that THC increases pregnenolone through activation of the CB<sub>1</sub> receptor.

Pregnenolone is synthesized in the mitochondria from cholesterol by cytochrome P450scc (16, 17). When high levels of steroid synthesis are needed (Fig. 2A), new cholesterol is provided first by hydrolysis of cytoplasmic cholesterol esters by the hormone-sensitive lipase (HSL), activated via phosphorylation of serine 660, and then by cholesterol transport into the mitochondria by the steroidogenic acute regulatory protein (StAR) (16, 17).

In Wistar rats, StAR proteins were not modified by THC administration (9 mg/kg) at any time (Fig. 2, C and D) (9). In contrast, THC induced a rapid (15 min after THC administration) increase in the levels of P450scc (Fig. 2, C and D) that was dependent on the activity of the extracellular signal-regulated kinases 1/2 mitogen-activated protein kinase (Erk1/2<sup>MAPK</sup>). This rapid increase in P450scc (9) was abolished by the selective inhibitor of Erk1/2<sup>MAPK</sup> phosphorylation SL327 (100 mg/kg; Fig. 2, C and D). Later (30 min after THC administration), an Erk1/2<sup>MAPK</sup>-independent mechanism sustained the increase in P450scc levels and phosphorylated HSL (Fig. 2, C and D). These coordinated modifications can contribute to the increase in pregnenolone induced by THC.


Among the behavioral and somatic CB<sub>1</sub> receptor-dependent effects of THC, the so-called “cannabinoid tetrad” (9, 18) is considered a prototypic signature of cannabinoid intoxication (14). The cannabinoid tetrad includes hypolocomotion, hypothermia, catalepsy, and analgesia. The administration of the inhibitor of pregnenolone synthesis aminoglutethimide (AMG, 50 mg/kg) (18) to C57BL/6N mice increased all of these behavioral and somatic effects of THC (Fig. 3, A to D). The injection of pregnenolone reversed the effects of AMG and inhibited the effects of THC but had no effect in animals that did not receive THC (Fig. 3, E and H). These data show that THC-induced production of pregnenolone exerts a negative feedback on CB<sub>1</sub> receptor activity.

Two well-known behavioral disturbances accompany cannabis use in humans: (i) an increase in food palatability and craving that can promote food intake (19, 20) and (ii) a decrease in memory performance (21). THC increases food intake in both sated rats and food-deprived C57BL/6N mice (22, 23) and also impairs memory consolidation in an object-recognition task in C57BL/6N mice (24). Pregnenolone administration (2 to 6 mg/kg) blocked THC-induced food intake in Wistar rats (Fig. 4A) and in C57BL/6N mice (Fig. 4B) and blunted the memory impairment induced by THC in mice (Fig. 4C), but it did not modify these behaviors per se (Fig. 4, A to C). As previously shown (24), THC-induced memory impairments were not due to nonspecific motor effects of THC, because THC did not significantly modify locomotor activity during the object-recognition task.

Cannabinoid drugs modulate brain activity and behavior principally by the activation of presynaptic CB<sub>1</sub> receptors, which inhibit the release of several neurotransmitters and in particular GABA and glutamate (25). We assessed the effect of THC on glutamate release (9) by measuring excitatory postsynaptic currents (EPSCs) in NAc principal neurons in brain slices obtained from adult Sprague-Dawley rats (Fig. 4, D and E). Bath application of THC (20  $\mu$ M) reliably inhibited synaptic transmission in control slices ( $34.3 \pm 3.7\%$  of inhibition). The effect of THC was significantly attenuated when slices were pre-treated with pregnenolone 100 nM ( $15.1 \pm 1.8\%$  of inhibition). These effects were probably due to a presynaptic action of pregnenolone. Thus, pregnenolone blocked the increase in paired-pulse ratio (PPR) induced by THC (9) but did not modify either the amplitude or the decay time of miniature EPSCs (mEPSCs) (table S2). Changes in PPR and in the mEPSC parameters studied here are indicators of changes in neurotransmitter release and in postsynaptic response, respectively.

To analyze the potential effects of pregnenolone on the development of cannabis abuse and dependence, we first studied dopamine release in the NAc (9). Activation of dopaminergic neurons and increase in NAc dopamine extracellular levels are common effects of most drugs of abuse and have been implicated in drug addiction (26).

In anesthetized Sprague-Dawley rats, we simultaneously used microdialysis and extracellular unitary recordings (9) to estimate dopamine release in the NAc and the firing activity of



**Fig. 3. The cannabinoid tetrad induced by THC was inhibited by pregnenolone.** In C57BL/6N mice, the administration, 30 min before THC, of AMG (50 mg/kg, ip), a P450scc inhibitor that blocks pregnenolone synthesis, increased the effects of THC: (A) hypolocomotion [ $F(3,98) = 13.8$ ,  $P < 0.001$ ], (B) hypothermia [ $F(3,98) = 4.7$ ,  $P < 0.01$ ], (C) catalepsy [ $F(3,98) = 2.1$ ,  $P < 0.05$ ], and (D) analgesia [ $F(3,98) = 2.2$ ,  $P < 0.05$ ]. (E to H) Pregnenolone [6 mg/kg, administered subcutaneously (sc)] administered at the same time as AMG (50 mg/kg, ip) prevented the increase in the responses to THC (10 mg/kg, ip) induced by AMG. Injection of pregnenolone (6 mg/kg, sc) alone 30 min before THC (10 mg/kg, ip) also reduced the behavioral effects of THC. Pregnenolone had no effects in animals that did not receive THC. Data are expressed as mean  $\pm$  SEM ( $n = 6$  to 12 animals per group). \* $P < 0.05$ ; \*\* $P < 0.01$ ; \*\*\* $P < 0.001$  as compared to vehicle-treated mice.