

## RESEARCH PAPER

# Blue light dose–responses of leaf photosynthesis, morphology, and chemical composition of *Cucumis sativus* grown under different combinations of red and blue light

Sander W. Hogewoning<sup>1,\*</sup>, Govert Trouwborst<sup>1</sup>, Hans Maljaars<sup>1</sup>, Hendrik Poorter<sup>2,3</sup>, Wim van Ieperen<sup>1</sup> and Jeremy Harbinson<sup>1</sup>

<sup>1</sup> Wageningen University, Department of Plant Sciences, Horticultural Supply Chains Group, Wageningen, The Netherlands

<sup>2</sup> ICG-3 (Phytosphäre), Forschungszentrum Jülich, D-52425 Jülich, Germany

<sup>3</sup> Plant Ecophysiology, Institute of Environmental Biology, PO Box 800.84, 3508 TB Utrecht, The Netherlands

\* To whom correspondence should be addressed. E-mail: [sander.hogewoning@gmail.com](mailto:sander.hogewoning@gmail.com)

Received 28 March 2010; Revised 27 April 2010; Accepted 27 April 2010

## Abstract

The blue part of the light spectrum has been associated with leaf characteristics which also develop under high irradiances. In this study blue light dose–response curves were made for the photosynthetic properties and related developmental characteristics of cucumber leaves that were grown at an equal irradiance under seven different combinations of red and blue light provided by light-emitting diodes. Only the leaves developed under red light alone (0% blue) displayed dysfunctional photosynthetic operation, characterized by a suboptimal and heterogeneously distributed dark-adapted  $F_v/F_m$ , a stomatal conductance unresponsive to irradiance, and a relatively low light-limited quantum yield for CO<sub>2</sub> fixation. Only 7% blue light was sufficient to prevent any overt dysfunctional photosynthesis, which can be considered a qualitatively blue light effect. The photosynthetic capacity ( $A_{\max}$ ) was twice as high for leaves grown at 7% blue compared with 0% blue, and continued to increase with increasing blue percentage during growth measured up to 50% blue. At 100% blue,  $A_{\max}$  was lower but photosynthetic functioning was normal. The increase in  $A_{\max}$  with blue percentage (0–50%) was associated with an increase in leaf mass per unit leaf area (LMA), nitrogen (N) content per area, chlorophyll (Chl) content per area, and stomatal conductance. Above 15% blue, the parameters  $A_{\max}$ , LMA, Chl content, photosynthetic N use efficiency, and the Chl:N ratio had a comparable relationship as reported for leaf responses to irradiance intensity. It is concluded that blue light during growth is qualitatively required for normal photosynthetic functioning and quantitatively mediates leaf responses resembling those to irradiance intensity.

**Key words:** Blue light, chlorophyll fluorescence imaging, cucumber (*Cucumis sativus*), dose–response curves, leaf mass per unit leaf area (LMA), light-emitting diodes (LEDs), photoinhibition, photosynthetic capacity, red light, starch accumulation.

## Introduction

Plant development and physiology are strongly influenced by the light spectrum of the growth environment. The underlying mechanisms of the effect of different growth spectra on plant development are not known in detail, although the involvement of photoreceptors has been demonstrated for a wide range of spectrum-dependent plant responses. Cryptochromes and phototropins are specifically blue light sensitive, whereas phytochromes are more

Abbreviations:  $A_{\max}$ , light-saturated assimilation;  $A_{\text{net}}$ , net assimilation; B, blue light percentage; Chl, chlorophyll;  $C_i$ ,  $C_a^{-1}$ , CO<sub>2</sub> concentration in leaf relative to CO<sub>2</sub> concentration in leaf chamber air; DW, dry weight; ETR, electron transport rate;  $F_v/F_m$ , ratio of variable to maximum fluorescence—the relative quantum efficiency for electron transport by PSII if all PSII reaction centres are open;  $g_{\text{sw}}$ , stomatal conductance;  $g_{\text{sw}}$  ratio, ratio of stomatal conductance on the adaxial and abaxial surface of the leaf; LED, light-emitting diode; LMA, leaf mass per unit leaf area; PNUE, photosynthetic nitrogen use efficiency; PSII, photosystem II; PSS, phytochrome photostationary state;  $R_{\text{dark}}$ , dark respiration;  $\alpha$ , light-limited quantum yield for CO<sub>2</sub> fixation;  $\Phi_{\text{PSII}}$ , relative quantum yield of PSII electron transport.

© 2010 The Author(s).

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/2.5>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

sensitive to red than to blue (Whitelam and Halliday, 2007). Blue light is involved in a wide range of plant processes such as phototropism, photomorphogenesis, stomatal opening, and leaf photosynthetic functioning (Whitelam and Halliday, 2007). At the chloroplast level, blue light has been associated with the expression of 'sun-type' characteristics such as a high photosynthetic capacity (Lichtenthaler *et al.*, 1980). Most studies assessing blue light effects on the leaf- or whole-plant level have either compared responses to a broad-band light source with responses to blue-deficient light (e.g. Britz and Sager, 1990; Matsuda *et al.*, 2008), or compared plants grown under blue or a combination of red and blue light with plants grown under red light alone (e.g. Brown *et al.*, 1995; Bukhov *et al.*, 1995; Yorio, 2001; Matsuda *et al.*, 2004; Ohashi *et al.*, 2006). Overall there is a trend to higher biomass production and photosynthetic capacity in a blue light-containing irradiance. Before the development of light-emitting diodes (LEDs) that were intense enough to be used for experimental plant cultivation (Tennessen *et al.*, 1994), light sources emitting wavelengths in a broader range than strictly the red (i.e. 600–700 nm) or blue (i.e. 400–500 nm) region were often used (e.g. Voskresenskaya *et al.*, 1977). Other wavelengths can interact with blue light responses. For example, green light has been reported to antagonize some blue light responses, such as stomatal opening and inhibition of hypocotyl elongation in seedlings (Folta and Maruhnich, 2007). The blue light enhancement effect on photosynthetic capacity appears to be greater when using combinations of red and blue light produced by LEDs than when broad-band light is made deficient in blue by a filter (e.g. for spinach compare Matsuda *et al.*, 2007 and 2008). This raises the question of whether plants exposed to red light alone suffer a spectral 'deficiency' syndrome, which may be reversed by blue light as well as by longer wavelengths.

Poorter *et al.* (2010) stress the importance of dose-response curves for quantitative analysis of the effects of environmental factors on plant phenotypes, allowing a better understanding of plant-environment interactions than the comparison of two treatments only. It is not clear whether the enhancement effect of blue light on leaf photosynthetic capacity is a qualitative threshold response or a quantitative progressive response, or a combination of both. Only few specific processes in leaves have been identified as quantitative blue light responses, such as chloroplast movement (Jarillo *et al.*, 2001) and stomatal conductance (Sharkey and Raschke, 1981). Matsuda *et al.* (2007) found a higher photosynthetic capacity for spinach leaves grown under  $300 \mu\text{mol m}^{-2} \text{s}^{-1}$  mixed red/blue irradiance containing  $30 \mu\text{mol m}^{-2} \text{s}^{-1}$  blue than for leaves grown under red alone. A higher blue light fraction did not yield a significant further enhancement in light-saturated assimilation ( $A_{\text{max}}$ ), which may be interpreted as a qualitative blue light effect. However, a quantitative blue light effect at quantum fluxes  $<30 \mu\text{mol m}^{-2} \text{s}^{-1}$  cannot be excluded.

A diverse choice of LEDs powerful enough for use as a growth irradiance source in controlled environments has

recently become available (e.g. Massa *et al.*, 2008). These LEDs allow the effect of light quality to be investigated independently of the amount of photosynthetic irradiance. LED illumination has been used here to study the response curves of a range of parameters related to leaf photosynthesis of plants that were grown at an irradiance with a proportion of blue light ranging from 0% to 100%. A range of other leaf characteristics important for the functioning of photosynthesis, such as stomatal development and behaviour, leaf mass per area (LMA), and the content of N, pigments, and carbohydrates, were also determined. The spectra and the extent of variation in the ratio of red and blue irradiance that can be achieved with LED lighting are dissimilar to field conditions. However, the responses of leaves to these unnatural environments provides the possibility to unravel the complex developmental and functional interactions that normally occur in the natural light environment.

## Materials and methods

### Plant material and growth conditions

Cucumber plants (*Cucumis sativus* cv. Hoffmann's Giganta) were sown in vermiculite and germinated under  $100 \mu\text{mol m}^{-2} \text{s}^{-1}$  cool white fluorescent lamps (TLD 50 W 840 HF, Philips, The Netherlands) in a climate chamber. After 1 week, when the cotyledons had just opened, the seedlings were transferred to a hydroponic system (Hoagland's solution,  $\text{pH}=5.9\pm0.2$ ;  $\text{EC}=1.2 \text{ mScm}^{-1}$ ) in a climate chamber. The day/night temperature was  $25^\circ\text{C}/23^\circ\text{C}$ , the relative humidity was 70%, and the  $\text{CO}_2$  concentration was ambient. All plants were subjected to  $100\pm5 \mu\text{mol m}^{-2} \text{s}^{-1}$  irradiance (16 h/8 h day/night) provided by a mixture of blue and red LEDs with dominant wavelengths of 450 nm and 638 nm, respectively (types Royal Blue and Red Luxeon K2, Lumileds Lighting Company, San Jose, CA, USA). The LEDs were equipped with lenses ( $6^\circ$  exit angle) and the arrays were suspended  $\sim 1$  m above the plants, so irradiance from the two LED types was well mixed. The lenses ensured that small differences in leaf height had only minor effects on the irradiance received. The seven different spectral treatments are expressed as the blue (B) light percentage: 0B, 7B, 15B, 22B, 30B, 50B, and 100B; the remaining percentage was red. Irradiance was measured routinely using a quantum sensor (LI-COR, Lincoln, NE, USA), but was also verified with a spectroradiometer (USB2000 spectrometer, Ocean Optics, Duiven, The Netherlands, calibrated against a standard light source). The difference in irradiance measured with the two devices was  $<2\%$  for the spectra used.

The plants were allowed to grow until the second leaf was fully mature (17–22 d after planting the seedlings) when it could be used for photosynthesis measurements. If necessary, the second leaf, which was the leaf used for all measurements, was supported in a horizontal position during growth to ensure that it received the specified irradiance.

### Stomata analysis

The stomatal conductance ( $g_{\text{sw}}$ ) was measured on three positions on each leaf surface using a leaf porometer (model SC-1, Decagon Devices, Inc., Pullman, WA, USA) prior to the gas exchange measurements (see below). The ratio of the average  $g_{\text{sw}}$  of the abaxial and adaxial leaf surface ( $g_{\text{sw}}$  ratio) was used in the calculations of the gas exchange parameters ( $n=6$ ). Additionally, silicon rubber impressions were made (see Smith *et al.*, 1989) on both the adaxial and abaxial surface of the leaves grown under 0B,

15B, 30B, and 50B ( $n \geq 3$ ). Stomatal density, length, and aperture were determined from images of the impressions using the procedure described in Nejad and van Meeteren (2005).

#### Leaf gas exchange and fluorescence measurements

Gas exchange and chlorophyll (Chl) fluorescence were measured using a custom-made leaf chamber within which 4.52 cm<sup>2</sup> of leaf surface was illuminated. A LI-7000 CO<sub>2</sub>/H<sub>2</sub>O gas analyser (LI-COR, Lincoln, NE, USA) measured the CO<sub>2</sub> and H<sub>2</sub>O exchange of the leaf and ambient atmospheric pressure. Leaf temperature was monitored by a thermocouple pressed against the abaxial leaf surface. A custom-made measuring-light source comprised of independently controllable red and blue LEDs with attached lenses, emitting a spectrum similar to that of the LEDs used for growth-light, was used to provide the required red/blue combination in the irradiance range 0–1700  $\mu\text{mol m}^{-2} \text{s}^{-1}$ . A polished steel reflector in the form of an inverted truncated cone (i.e. the inlet to the reflector was larger than the outlet) allowed the irradiance to be well mixed and equally distributed over the leaf surface. The gas mix used contained 380  $\mu\text{mol mol}^{-1}$  CO<sub>2</sub>, 20.8 $\pm$ 0.4 mmol mol<sup>-1</sup> H<sub>2</sub>O, and either 210 mmol mol<sup>-1</sup> or 20 mmol mol<sup>-1</sup> O<sub>2</sub> (ambient O<sub>2</sub> or low O<sub>2</sub>), dependent on the type of measurement. A flow rate of 200–700 ml min<sup>-1</sup> was used, depending on the CO<sub>2</sub> depletion which ranged from 18  $\mu\text{mol mol}^{-1}$  to 26  $\mu\text{mol mol}^{-1}$  at saturating irradiance. The equations developed by von Caemmerer and Farquhar (1981) were used to calculate assimilation,  $g_{\text{sw}}$ , and the CO<sub>2</sub> concentration in the substomatal cavity of the leaf relative to that in the leaf chamber air ( $C_i/C_a$ ) from the gas exchange data. The boundary layer resistance of both leaf surfaces in the leaf chamber during gas exchange measurements was estimated using the method of Jarvis (1971). Chl fluorescence was measured using a PAM 101 Chl fluorometer with an emitter detector unit (model 101 ED; Heinz Walz, Effeltrich, Germany). The modulated red measuring-light intensity was <0.5  $\mu\text{mol m}^{-2} \text{s}^{-1}$ . A 250 W quartz-halogen lamp connected to an additional optical fibre provided a saturating light pulse (7500  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ) to allow measurement of the  $F_m$  or  $F_m'$  relative fluorescence yield (Baker *et al.*, 2007). The fibres were fixed ~4 cm above the leaf chamber at such an angle that they did not interfere with the actinic light beam.

Irradiance–response curves were measured on fully expanded second leaves, and each growth-light treatment was performed twice. As there were no significant differences between the two repetitions, the individual plants from the two repetitions were treated as independent repetitions ( $n=6$ ) in the analysis. An ambient O<sub>2</sub> concentration was used for these measurements. After clamping a leaf in the leaf chamber, it was dark adapted for 30 min, and dark respiration ( $R_{\text{dark}}$ ) and the dark-adapted  $F_v/F_m$  (Baker *et al.*, 2007) were measured. The irradiance–response curve was measured using a spectrum identical to that under which the plants were grown, using 14 intensities in the range 0–1700  $\mu\text{mol m}^{-2} \text{s}^{-1}$ . The leaves were subjected to each irradiance for at least 20 min, when steady-state assimilation was amply reached. The highest irradiances were omitted if CO<sub>2</sub> fixation clearly became light-saturated at lower irradiances. At an irradiance of 100  $\mu\text{mol m}^{-2} \text{s}^{-1}$ , which is equal to the irradiance during growth, the relative quantum yield of photosystem II (PSII) electron transport ( $\Phi_{\text{PSII}}$ ) was measured using the method of Genty *et al.* (1989). After measuring the irradiance–response curve, the plant was left overnight in the dark in a climate room and the following day samples were taken from the measured leaf in order to measure the light absorbance spectrum, leaf mass per area (LMA), and pigment- and N-content (see below).

In order to assess the possibility that  $C_i$  was limiting assimilation at low irradiance, the relationship between assimilation and electron transport rate (ETR) was investigated in more detail. Under photorespiratory conditions a lower assimilation per unit ETR is expected for a leaf with a  $C_i$  that is limiting for assimilation than for a leaf with no limiting  $C_i$ . Under non-photorespiratory

conditions no difference is to be expected (Harbinson *et al.*, 1990). Additional gas exchange and fluorescence measurements were made on leaves grown under 0B and 30B using seven different incident irradiances (0–100  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ) and both ambient and low O<sub>2</sub> ( $n=3$ ). Chl fluorescence measurements were made at each irradiance to determine  $\Phi_{\text{PSII}}$  once CO<sub>2</sub> fixation had stabilized, after which the actinic irradiance was switched off to measure  $R_{\text{dark}}$ . Gross assimilation ( $A_{\text{gross}}$ ) was calculated as net assimilation ( $A_{\text{net}}$ ) plus  $R_{\text{dark}}$ , which assumes, as is commonly done, that  $R_{\text{dark}}$  is a reasonable estimate of respiration in the light. Light absorbance (see below) was measured directly after measuring the photosynthesis irradiance–response. The product of the absorbed actinic irradiance and  $\Phi_{\text{PSII}}$  serves as an index for ETR (e.g. Kingston-Smith *et al.*, 1997). The distribution of dark-adapted  $F_v/F_m$  over these 0B- and 30B-grown leaves was measured by means of Chl fluorescence images. Images of three different leaves from each treatment were made using a PSI Fluorcam 700MF Chl fluorescence imaging system (PSI, Brno, Czech Republic), using the procedure described in Hogewoning and Harbinson (2007).

#### Measurement of leaf light absorbance

Leaf light absorbance was calculated in 1 nm steps in the range 400–800 nm from measurements of leaf reflectance and transmittance made on 12 leaf discs per leaf. Details of the procedure and measurement system, which consisted of two integrating spheres, each connected to a spectrometer and a custom-made light source, are described in Hogewoning *et al.* (2010) and Zheng *et al.* (2010). The integrated absorbance of the actinic measuring irradiance used during gas exchange measurements was subsequently calculated by multiplying the relative leaf absorbance spectrum by the spectrum of the measuring-light.

#### LMA, nitrogen, pigment, and carbohydrate analysis

From each leaf, 10 leaf discs (1.28 cm<sup>2</sup>) were cut randomly over the leaf area, avoiding the leaf margins and main veins. The discs were stored at –22 °C, freeze dried, and weighed, and LMA was calculated. After weighing, the C and N contents were determined for all treatments by a C/N analyser ( $n=5$ ) and the nitrate content was determined for the treatments 0B and 30B ( $n=4$ ) according to Trouwborst *et al.* (2010).

An additional eight leaf discs (0.65 cm<sup>2</sup>) were cut from the same leaf and stored in 10 ml of dimethylformamide (DMF) in the dark at –22 °C. The absorbance of the extract was measured in the range 400–750 nm using a Cary 4000 spectrophotometer (Varian Instruments, Walnut Creek, CA, USA), and the Chl and carotenoid concentrations were calculated using the equations of Wellburn (1994).

The carbohydrate content of leaves grown under 0B, 30B, and 100B was measured by cutting 10–15 discs (1.28 cm<sup>2</sup>) from one side of the main vein at the end of the photoperiod and 10–15 discs from the other side of the main vein just before the start of the photoperiod ( $n=4$ ). Soluble carbohydrate and starch concentrations were analysed as described in Hogewoning and Harbinson (2007).

#### Curve fitting and statistics

The photosynthesis data measured to obtain light–response curves of the leaves grown under different blue/red combinations were fitted with a non-rectangular hyperbola (Thornley, 1976) using the non-linear fitting procedure NLIN in SAS (SAS Institute Inc. 9.1, Cary, NC, USA) in order to determine the light-limited quantum yield for CO<sub>2</sub> fixation ( $\alpha$ ).

Tukey's HSD was used to make post-hoc multiple comparisons among spectral treatment means from significant one-way analysis of variance (ANOVA) tests ( $P < 0.05$ ), and regression analysis was used to test for significant differences ( $P < 0.05$ ) between the slope of the  $A_{\text{gross}} - \Phi_{\text{PSII}} \times \text{absorbed measuring-light}$  relationship




using Genstat (release 9.2, Rothamsted Experimental Station, Harpenden, UK).

Results

Leaf photosynthesis

The  $A_{\max}$  differed significantly for the leaves grown under different blue (B) light percentages (Fig. 1). Increasing the blue light fraction from 0% to 50% resulted in an increasing  $A_{\max}$ , with the greatest increase occurring at the increase from 0% to 7% blue. The 100B-grown leaves had an  $A_{\max}$  that was lower than that of the 50B leaves. The light-limited quantum yield for CO<sub>2</sub> fixation ( $\alpha$ ) was lowest for 0B and 100B leaves and highest for the 7B–30B leaves (within this range there was no significant difference in  $\alpha$ ; Table 1). Dark respiration was lowest for 0B leaves and tended to increase with blue light percentage, except for 100B (Table 1), similar to the pattern found for  $A_{\max}$ . The dark-adapted  $F_v/F_m$  was typical for an unstressed leaf (i.e.  $\geq 0.8$ ) in all treatments, except 0B, where it was significantly reduced (Table 1). The  $\Phi_{PSII}$  measured at growth-light intensity (i.e. 100  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ) and spectrum was similar



**Fig. 1.** The effect of light quality (the proportion of total PAR that is from the blue rather than from the red part of the spectrum) during growth on the photosynthetic capacity ( $A_{\max}$ ) of cucumber leaves. Error bars indicate the SEM ( $n=6$ ).

**Table 1.** Different parameters measured or calculated on leaves grown under different light qualities (the proportion of total PAR that is from the blue rather than from the red part of the spectrum).

Different letters indicate significant differences ( $P \leq 0.05$ ;  $n=5$  or  $n=6$ , no variation for PSS).

Blue light percentage	0	7	15	22	30	50	100
$F_v/F_m$	0.76 b	0.80 a	0.80 a	0.80 a	0.81 a	0.81 a	0.81 a
$\Phi_{PSII}$	0.65 d	0.74 c	0.76 b	0.76 a,b	0.76 a, b	0.77 a	0.76 a,b,c
$F_v/F_m - \Phi_{PSII}$	0.110 a	0.055 b	0.044 c	0.040 c	0.042 c	0.034 c	0.044 b,c
Quantum yield CO <sub>2</sub> fixation ( $\alpha$ )	0.045 c	0.052 a,b	0.053 a	0.053 a	0.053 a	0.048 b,c	0.045 c
$R_{\text{dark}}$ ( $\mu\text{mol m}^{-2} \text{s}^{-1}$ )	0.93 d	1.17 c	1.29 a,b,c	1.39 a,b	1.27 b,c	1.45 a	1.33 a,b,c
$g_{\text{sw}}$ ratio (abaxial:adaxial)	2.7 a	2.6 a	2.1 a,b	1.7 b,c	1.7 b,c	1.4 c	1.7 b,c
Integrated absorbance	90.0 d	92.1 c	92.4 b,c	93.1 b,c	94.0 a,b	93.7 b	95.4 a
Chl a:b ( $\text{g g}^{-1}$ )	3.24 d	3.36 c	3.51 a,b	3.48 a,b	3.42 b,c	3.54 a	3.54 a
N (% DW)	5.7 a	6.0 a	5.7 a	6.0 a	6.1 a	6.0 a	6.2 a
C (% DW)	39.6 a	38.0 a	36.8 a	38.7 a	37.7 a	37.6 a	37.7 a
C:N ( $\text{g g}^{-1}$ )	6.9 a	6.4 a,b	6.5 a,b	6.4 a,b	6.2 b	6.2 b	6.1 b
Chl:N ( $\text{g g}^{-1}$ )	5.1 a	4.3 b,c	4.6 a,b	4.1 b,c,d	4.3 b,c	3.9 c,d	3.7 d
PSS (phytochromes)	0.89	0.89	0.89	0.89	0.88	0.87	0.51


for the 15B–100B leaves, but was markedly lower for 0B leaves and slightly, but significantly, lower for 7B leaves.

Concerning the more detailed measurements of the photosynthesis irradiance–response between 0  $\mu\text{mol m}^{-2} \text{s}^{-1}$  and 100  $\mu\text{mol m}^{-2} \text{s}^{-1}$  incident irradiance on 0B- and 30B-grown leaves,  $A_{\text{gross}}$  was markedly higher for the low O<sub>2</sub> measurements than it was for the ambient O<sub>2</sub> measurements (Fig. 2). At all light intensities,  $\Phi_{PSII}$  was consistently lower for the 0B leaves than it was for the 30B leaves. In both treatments the O<sub>2</sub> concentration did not affect  $\Phi_{PSII}$  (not shown). The absorbance in the green region of the spectrum was 5–10% lower for the 0B- and 100B-grown leaves than for the other treatments, whereas differences were negligible for the blue and red region (not shown). Only the red and blue wavelength regions are relevant for integrated absorbed irradiance in this experiment. The integrated absorbance of the growth and measuring-light increased with the percentage of blue light (Table 1), as the blue light was better absorbed than the red light. At both low and ambient O<sub>2</sub> concentration there were no significant differences between 0B and 30B for the linear regression between  $A_{\text{gross}}$  and the product of  $\Phi_{PSII}$  and the absorbed actinic irradiance (Fig. 2).


The images of dark-adapted  $F_v/F_m$  obtained via Chl fluorescence imaging showed conspicuous differences between the 0B and 30B leaves. Whereas the images from 30B-grown leaves were perfectly homogeneous with an  $F_v/F_m > 0.8$ , the images of the 0B-grown leaves showed a heterogeneous distribution with dark-adapted  $F_v/F_m$  values of  $\sim 0.8$  adjacent to the veins and with zones of lower  $F_v/F_m$  (typically 0.55–0.70) between the veins (Fig. 3). The 0B leaves also occasionally appeared slightly chlorotic between the veins.

Stomatal effects

There was a considerable stomatal conductance ( $g_{\text{sw}}$ ) calculated from gas exchange data in the dark-adapted state (Fig. 4B). As the photoperiod of the plants in their growth




**Fig. 2.** Relationship between gross  $\text{CO}_2$  assimilation ( $A_{\text{gross}}$ ) and the product of  $\Phi_{\text{PSII}}$  and the actinic measuring-light absorbed by the leaves, which serves as an index of electron transport (e.g. Kingston-Smith *et al.*, 1997), at an incident irradiance  $\leq 100 \mu\text{mol m}^{-2} \text{s}^{-1}$ . The cucumber leaves were grown under and also measured with 0B (=100% red; circles) and 30B (squares) irradiance, and gas exchange was measured under low (open symbols) and ambient  $\text{O}_2$  (filled symbols). Gross assimilation was calculated as dark respiration plus net assimilation. The slopes of the regression lines are significantly different for the two  $\text{O}_2$  levels ( $P < 0.001$ ), but not for the spectral treatments ( $P \geq 0.23$ ).



**Fig. 3.** Image of the dark-adapted  $F_v/F_m$  distribution over an 0B (=100% red; A) and 30B (B) irradiance-grown cucumber leaf. The mixed blue-red-grown leaf (B) has a homogeneous  $F_v/F_m$  distribution centred around an  $F_v/F_m$  of 0.82, whereas the 0B-grown leaf (A) has a heterogeneous distribution with a high  $F_v/F_m$  around the veins and lower values between the veins.

environment started 1 h before leaves were dark adapted in the leaf chamber, the absence of complete stomatal closure may be due to the diurnal rhythm of the stomata. Also, a significant night-time  $g_{\text{sw}}$  is not unusual, especially for leaves with a high daytime  $g_{\text{sw}}$  (Snyder *et al.*, 2003), such as cucumber. Moreover, a substantial night-time  $g_{\text{sw}}$  has been reported to occur in many horticultural species, and ample water availability (e.g. hydroponics as used here) can increase night-time  $g_{\text{sw}}$  (Caird *et al.*, 2007). The  $g_{\text{sw}}$  of leaves grown and measured using 0B was lowest of all the treatments and did not respond to increases in measuring irradiance intensity. Even using 30B or 100B as a measuring irradiance spectrum on the 0B-grown leaves at either  $100 \mu\text{mol m}^{-2} \text{s}^{-1}$  irradiance or saturating irradiance had




**Fig. 4.** Response of net assimilation ( $A_{\text{net}}$ ; A), stomatal conductance ( $g_{\text{sw}}$ ; B), and leaf internal  $\text{CO}_2$  concentration relative to that of the leaf chamber air ( $C_i/C_a$ ; C) to irradiance for cucumber leaves grown under different light qualities (the proportion of total PAR that is from the blue rather than from the red part of the spectrum). The actinic light quality was identical to that during growth. Error bars indicate the SEM ( $n=6$ ).

no effect on their  $g_{\text{sw}}$  (data not shown). In all other treatments,  $g_{\text{sw}}$  increased with increasing irradiance ( $>100 \mu\text{mol m}^{-2} \text{s}^{-1}$ ). Consistent with the low and constant  $g_{\text{sw}}$ , the  $C_i/C_a$  of the 0B-grown leaves decreased more with increasing irradiance than that of the other treatments (Fig. 4C). Data of  $g_{\text{sw}}$  and  $C_i/C_a$  for the 30B and 100B leaves are not shown in Fig. 4 due to instrument failure.


The  $g_{\text{sw}}$  measured using a porometer also increased with increasing blue light in the growth spectrum (not shown). The ratio of  $g_{\text{sw}}$  on the abaxial and adaxial leaf surface ( $g_{\text{sw}}$  ratio) became smaller with an increasing percentage of blue light (Table 1). The stomatal counts on both leaf sides paralleled these results, as the number of stomata on the adaxial leaf surface significantly increased with increasing blue percentage, whereas on the abaxial leaf surface no significant changes were found (not shown), resulting in a decreasing stomatal ratio with increasing blue light (Fig. 5). No significant changes in stomatal length and guard cell width were found for the different treatments (not shown).

## LMA and nitrogen, pigment, and carbohydrate content

The LMA increased with increasing percentage of blue up to 50% (Fig. 6A). Similar to the  $A_{\max}$ –blue percentage relation-



**Fig. 5.** Ratio of stomatal density (open bars;  $n \geq 3$ ) and stomatal conductance measured with a porometer (filled bars;  $n=6$ ) for the abaxial and adaxial leaf surface of cucumber leaves grown under different light qualities (the proportion of total PAR that is from the blue rather than from the red part of the spectrum; both parameters are labelled 'stomatal ratio' in the plot). Error bars indicate the SEM and letters indicate significant differences ( $P \leq 0.05$ ). No significant differences between the individual means of the stomatal density ratio were found; however, the linear component of the stomatal density ratio–blue light percentage relationship was significant ( $P=0.04$ ). The decrease in stomatal density ratio with increasing blue light percentage was due to an increasing stomatal density on the adaxial leaf surface.



**Fig. 6.** The effect of light quality (the proportion of total PAR that is from the blue rather than from the red part of the spectrum) during growth on the chlorophyll content per unit leaf area (A, filled symbols, left axis), leaf mass per unit leaf area (LMA; A, open symbols, right axis), and the percentage chlorophyll in the leaf on a dry weight basis (B, squares).

ship (Fig 1), the increase in LMA was relatively greatest when the growth irradiance was changed from 0B to 7B. The total Chl content (Chl *a*+Chl *b*; Fig. 6A) and total carotenoid content (not shown) per unit leaf area increased in a similar way to LMA, increasing with percentage blue up to 50%. The Chl *a*:*b* ratio was significantly lower for 0B and 7B than at higher blue percentages (Table 1).

Leaf N content and C content per unit dry weight (DW) did not differ significantly between the treatments (Table 1). When expressed per unit leaf area the N and C content therefore depended on the percentage blue light in a way that was similar to LMA (Fig. 6A). The C:N ratio, however, was significantly higher for the 0B treatment than it was for the 30B, 50B, and 100B treatments. The nitrate part of total leaf N was not significantly different for the 0B and 30B leaves and was only 8.8% and 6.4%, respectively.

Chl content per unit leaf area correlates well with LMA (Fig. 6A), though there is a small but significant decrease in the Chl content per unit leaf DW as the percentage blue light in the growth irradiance increases (Fig. 6B). For all treatments  $A_{\max}$  correlated positively with LMA and Chl content per area leaf, except for Chl content of the 100B leaves (Fig. 7). With an increasing percentage blue light during growth,  $A_{\max}$  per unit Chl increases up to 22B, whereas at higher percentages of blue there are no differences between the treatments (Fig. 8A). A similar pattern can be seen for  $A_{\max}$  per unit leaf DW (Fig. 8A) and  $A_{\max}$  per unit N, which is the photosynthetic N use efficiency (PNUE; Fig. 8B). On a DW basis, the Chl:N ratio decreases significantly with increasing percentage blue (Table 1).


The leaf carbohydrate content (on a unit weight basis) was negligibly low at the end of the night period for all treatments (Table 2). At the end of the photoperiod, a considerable amount of carbohydrates, which were mainly comprised of starch and smaller quantities of sucrose, was present in the leaves, with the highest values in the leaves grown under 30B.

## Discussion


Peculiarly, whereas parameters such as  $A_{\max}$ , leaf composition, and LMA depended on the percentage of blue light during growth, only the leaves that developed under 0B (100% red light) had a suboptimal  $F_v/F_m$ , a low light-limited quantum efficiency for CO<sub>2</sub> fixation ( $\alpha$ ; Table 1), and a stomatal conductance ( $g_{sw}$ ) that was unresponsive to irradiance (Fig. 4). Such effects on leaves have, to the best of our knowledge, not been reported before and highlight the fundamental difference between leaf adaptation to the growth spectrum and the instantaneous spectral effect on photosynthesis. Instantaneous photosynthetic rates are relatively high when a leaf is illuminated with red light (e.g. McCree, 1972; Inada, 1976).

### Disorders in leaf physiology associated with growth under red light alone

A lower photosynthetic rate in plants grown under red light alone has been shown for several crop plants. Matsuda



**Fig. 7.** Relationship of leaf photosynthetic capacity ( $A_{\max}$ ) to leaf mass per unit leaf area (A) and chlorophyll content per unit leaf area (B) of cucumber grown under different combinations of red and blue light at an equal irradiance. The order of the values related to the data points corresponds to the blue light percentage under which the leaves were grown, except for the encircled data point which refers to the 100% blue treatment.



**Fig. 8.** The effect of light quality (the proportion of total PAR that is from the blue rather than from the red part of the spectrum) during growth of cucumber on leaf photosynthetic capacity ( $A_{\max}$ ) reached per unit chlorophyll (A, filled symbols, left axis), per unit leaf dry weight (A, open symbols, right axis), and per unit N (B, squares).

*et al.* (2004) found a lower photosynthetic rate for rice grown under red LEDs alone than for plants grown under a mixture of red and blue LEDs. Similar results were found for wheat (Goins *et al.*, 1997), which had a lower photosynthesis and DW accumulation when grown under red alone compared with growth under white fluorescent tubes or under red light supplemented with blue. While Yorio *et al.* (2001) reported a lower DW accumulation in radish, spinach and lettuce grown under red LEDs alone than under white fluorescent tubes or red supplemented with

**Table 2.** Carbohydrate content (mg g<sup>-1</sup> DW) of leaves grown under different light qualities (the proportion of total PAR that is from the blue rather than from the red part of the spectrum). Different letters indicate significant differences ( $P \leq 0.05$ ;  $n=4$ ).

Blue %	End dark period			End photoperiod		
	0	30	100	0	30	100
Glucose	0.4 a	0.2 a	0.4 a	0.5 a	0.4 a	0.4 a
Sucrose	0.5 a	0.3 a	0.4 a	8.4 b	9.6 b	13.2 a
Starch	1.1 a	0.6 a	0.8 a	45.1 b	55.8 a	39.5 b

blue, only radish developed a lower photosynthetic rate when grown under red LEDs (as we also found for cucumber; Figs 1, 4A). This suggests that vulnerability to decreases in photosynthetic rate associated with growth under red light alone may be subject to genetic variation.

The low  $A_{\max}$  of the leaves that developed under 0B (Fig. 1) cannot be attributed to a low leaf N content, as the PNUE at  $A_{\max}$  is lower for the 0B treatment than for the other treatments (Fig. 8B). Chl content and LMA can also be ruled out, as  $A_{\max}$  expressed per unit leaf DW and per unit Chl is also lower for the 0B leaves (Fig. 8A). The nitrate fraction of the leaf N content has been reported to be relatively higher in leaves grown under low irradiance than those grown under a high irradiance (e.g. Felipe, *et al.*, 1975). In the present study this nitrate effect on PNUE can be excluded as in both in the 0B and 30B leaves N in the form of nitrate was <10% of the total N content. The unresponsiveness of the stomata of 0B-grown leaves did limit  $A_{\max}$  due to a more restricted CO<sub>2</sub> diffusion into the leaf, as reflected by the lower  $C_i$   $C_a^{-1}$  with increasing measuring irradiance in the 0B leaves compared with the other treatments (Fig. 4).

In contrast to  $A_{\max}$ , the low  $\alpha$  found for the 0B treatment (Table 1) is entirely related to a lower  $\Phi_{PSII}$  and not to a low  $C_i$  due to a low  $g_{sw}$  (Fig. 4), as under both ambient O<sub>2</sub> and non-photorespiratory conditions the relationship between  $A_{gross}$  and an index of ETR (the product of  $\Phi_{PSII}$  and absorbed irradiance) did not differ significantly for the 0B



and the 30B leaves (Fig. 2). If  $C_i$  was limiting assimilation of the 0B leaves at low irradiance,  $A_{\text{gross}}$  per unit ETR would have been lower for 0B than for 30B at ambient  $O_2$  but not at low  $O_2$  (e.g. Harbinson *et al.*, 1990). Therefore, the underlying cause of the relatively low photosynthetic rates at low irradiance of the 0B-grown leaves may be due to disorders in the development and functioning of the photosynthetic machinery itself. During the photosynthesis measurements the measuring-light spectrum was identical to the growth-light, so a higher  $\alpha$  would be expected for the 0B treatment as the quantum yield for incident red light is known to be higher than that of blue light (McCree, 1972; Inada, 1976). Where the relatively low  $\alpha$  measured for the treatments containing a high blue light percentage (50B, 100B) was to be expected based on the differences in quantum yields for the different wavelengths, the low  $\alpha$  for the 0B treatment is unexpected and points to problems in the development and operation of photosynthesis. An  $F_v/F_m$  below 0.8, as measured for the 0B leaves, is normally associated with damage or long-term down-regulation of PSII in response to stress (e.g. Baker, 2008). Evidently red light alone, or the absence of blue light during growth, results in a dysfunction of the photosynthetic machinery with a particularly adverse effect on leaf tissue regions between the veins (Fig. 3). Matsuda *et al.* (2008) reported an  $F_v/F_m \geq 0.8$  for spinach leaves grown under white fluorescent light deficient in blue, so wavelengths beyond the blue region may also prevent a loss of  $F_v/F_m$ , as found for 100% red in this study.

Several diverse, spectrally related factors have been associated with inhibition of photosynthesis. Feedback down-regulation of photosynthesis is associated with carbohydrate accumulation in leaves (e.g. Stitt, 1991; Paul and Foyer, 2001). Britz and Sager (1990) found lower leaf photosynthesis associated with higher starch content at the end of the night period in soybean and sorghum leaves grown under low pressure sodium lamps emitting very little blue light and mainly amber/red light ( $\sim 595$  nm), compared with leaves grown under daylight fluorescent tubes. In the case of the present experiments any such effects on carbohydrate transport and metabolism can be discounted as no differences in carbohydrate content at the end of the dark period were found between the treatments (Table 2). In wheat seedlings, inhibition of PSI and PSII development and Chl synthesis was reported upon exposing the root-shoot transition zone to  $500 \mu\text{mol m}^{-2} \text{s}^{-1}$  pure red light (Sood *et al.*, 2004), suggesting an unidentified problem related to transport of substances within the plant. In the present experiment, Chl content on a leaf DW basis was not impaired in the 0B treatment (Fig. 6); however, the higher  $F_v/F_m$  adjacent to the veins (Fig. 3) and the occasional chlorotic appearance between the veins also point to a potential transport problem. Schmid and co-workers related a depressed  $F_v/F_m$  and photosynthesis in chloroplasts of red light-grown green algae *Acetabularia* to uncoupling of antennae and PSII reaction centres due to reduced amounts of core antenna Chl-protein complexes (Wenicke and Schmid, 1987; Schmid *et al.* 1990a, b). The involvement of a blue light/UV-A photosensory pathway

in the maintenance of PSII core protein synthesis has been postulated by Christopher and Mullet (1994), and Mochizuki *et al.* (2004) found a threshold intensity of  $5 \mu\text{mol m}^{-2} \text{s}^{-1}$  blue light (470 nm) for activation of the PSII core protein D2-encoding gene *psbD* in *Arabidopsis* acting via cryptochromes, along with a non-blue-specific activation signal. An impaired ability to synthesize core proteins may be related to the low  $F_v/F_m$  and  $\alpha$  that were found for the 0B-grown cucumber leaves; however, this theory cannot be directly linked to a problem with transport within the plant, as indicated by the heterogeneous  $F_v/F_m$ .

### Blue light dose-responses

The physiological disorders associated with leaf development under red light alone were eliminated by adding only a small amount of blue light (7% or  $7 \mu\text{mol m}^{-2} \text{s}^{-1}$ ; Fig. 1). Beside this response to blue, which may be characterized as a 'qualitative' or 'threshold' effect, the increase in  $A_{\text{max}}$  upon increasing the blue light percentage up to 50B clearly indicates that leaf photosynthesis also responds quantitatively to blue light.

The quantitative increase in  $A_{\text{max}}$  with an increasing proportion of blue light was associated with an increase in LMA (Fig. 7A), Chl content, and N per unit area (Table 1; Fig. 7B) and  $g_{\text{sw}}$  at saturating irradiance (Fig. 4B). The larger  $g_{\text{sw}}$  is due both to a larger number of adaxial stomata (Fig. 5) and a greater stomatal aperture. Blue light deficiency has been associated with a lower LMA in soybean (Britz and Sager, 1990), consistent with the lowest LMA that was found for the 0B-grown leaves here. A higher irradiance is usually found to lead to both a higher LMA and  $A_{\text{max}}$  (Poorter *et al.*, 2009). The present results show that the quantitative relationship between LMA and  $A_{\text{max}}$  with increasing irradiance (Poorter *et al.*, 2009, 2010) is also found for a varying blue percentage at a constant irradiance (Fig. 7A). In general, in parallel with leaf responses to irradiance, blue light is shown to stimulate 'sun-type' characteristics on the leaf level, even at the relatively low growth irradiance used in this study.

The question remains of which blue light-regulated response(s) can explain the differences in  $A_{\text{max}}$  of leaves grown under different blue light percentages? At a blue light percentage  $\geq 22\%$   $A_{\text{max}}$  appears to change proportionally to changes in LMA, Chl, and PNUE (Fig. 8), although Chl per leaf DW (Fig. 6B) and Chl:N (Table 1) decrease slightly with an increasing percentage of blue light. Similar relationships between these leaf traits are usually observed with increasing irradiances, where  $A_{\text{max}}$  increases proportionally with LMA and N content per unit leaf area, and Chl:N decreases (e.g. Evans and Poorter, 2001). Leaf N content may therefore indeed be a limiting factor for  $A_{\text{max}}$  of leaves grown at an irradiance  $\geq 22\text{B}$ . Regulation of potential  $A_{\text{max}}$  due to restrictions in cell size and the number of cell layers in a mature leaf as proposed by Oguchi *et al.* (2003) is also well in line with the correlation found between LMA and  $A_{\text{max}}$  in the present experiment. A restriction in intercellular space per unit leaf area may be expected to be associated



with a limitation of N-requiring components of the photosynthetic machinery per unit leaf area. More unusual is the lower  $A_{\max}$  per unit LMA, Chl, and N found for leaves grown under an irradiance containing  $\leq 15\text{B}$  (Fig. 8). These results indicate that cell space within the leaf, N availability, and pigment content were sufficiently large to allow a higher  $A_{\max}$ . Hogewoning *et al.* (2010) likewise found a lower  $A_{\max}$  per unit LMA for cucumber leaves grown under high pressure sodium light (5% blue) compared with leaves grown under fluorescent tubes (23% blue) and an artificial solar spectrum (18% blue). Apparently leaves grown at an irradiance containing  $\leq 15\text{B}$  are subject to limitations which may be related to the disorders associated with 0B leaves as discussed above, whereas at  $\geq 22\text{B}$  the relationships between  $A_{\max}$  and LMA, N, and Chl are very similar to usual leaf responses to irradiance.

The Chl *a:b* ratio was also conspicuously lower for 0B and 7B leaves, but remained stable at  $>15\text{B}$  (Table 1). This response is not in accordance with the usually measured increasing Chl *a:b* ratio with increasing irradiance during growth (Evans and Poorter, 2001), in contrast to the responses of the other leaf traits measured, which are in accordance with usual responses to irradiance.

#### Leaf responses to growth under blue light alone

Though the responses of  $A_{\max}$  (Fig. 1), LMA, and Chl content (Fig. 6A) in the range 0B–50B display clear progressive trends, the results for the 100B treatment deviate from those trends. In contrast to 0B, 100B leaves did not show any signs of dysfunctional photosynthesis. One conspicuous contrast between red and blue light is the absence of cryptochrome and phototropin stimulation in pure red, whereas pure blue does stimulate cryptochromes, phototropins, and also phytochromes (Whitelam and Halliday, 2007). The 100B leaves invested relatively little in Chl considering their  $A_{\max}$  (Fig. 7). The relative amount of active phytochrome expressed as the phytochrome photostationary state (PSS; calculated according to Sager *et al.*, 1988) of the 100B leaves is also markedly lower than that of the other red/blue combinations (Table 1), which may indicate a role for phytochrome activity in the regulation of the Chl content– $A_{\max}$  relationship. As LMA has been shown to be much less affected than  $A_{\max}$  at spectra containing relatively little blue (Fig. 8A; high pressure sodium light-grown leaves in Hogewoning *et al.*, 2010), the lower  $A_{\max}$  of 100B leaves compared with 50B leaves may be related to a limitation in LMA due to the absence of responses regulated by red light.

#### Conclusions

In this study, blue light has been shown to trigger both a qualitative signalling effect enabling normal photosynthetic functioning of cucumber leaves and a quantitative response stimulating leaf development normally associated with acclimation to irradiance intensity. Leaf acclimation to irradiance intensity may therefore be regulated by a limited range of wavelengths instead of the full PAR spectrum. Varying the blue light fraction offers the possibility to

manipulate leaf properties under a low irradiance such that they would normally be associated with high irradiances. The possibility to grow plants under relatively low irradiance in a plant growth facility, with a relatively high photosynthetic capacity able to withstand irradiances under field conditions, is a useful practical consequence for research and agriculture.

#### Acknowledgements

This research is supported by the Dutch Technology Foundation STW, Applied Science Division of the NWO, and the Technology Program of the Ministry of Economic Affairs, Philips, and Plant Dynamics BV. We are grateful to Joost Ruijsch, Evert Janssen, and Gradus Leenders (equipment development), Annie van Gelder and Joke Oosterkamp (stomata analysis), Arjen van de Peppel (biochemical analysis), and Hennie Halm C/N measurements).

#### References

- Baker NR. 2008. Chlorophyll fluorescence: a probe of photosynthesis *in vivo*. *Annual Review of Plant Biology* **59**, 89–113.
- Baker NR, Harbinson J, Kramer DM. 2007. Determining the limitations and regulation of photosynthetic energy transduction in leaves. *Plant, Cell and Environment* **30**, 1107–1125.
- Britz SJ, Sager JC. 1990. Photomorphogenesis and photoassimilation in soybean and sorghum grown under broad-spectrum or blue-deficient light-sources. *Plant Physiology* **94**, 448–454.
- Brown CS, Schuerger AC, Sager JC. 1995. Growth and photomorphogenesis of pepper plants under red light-emitting diodes with supplemental blue or far-red lighting. *Journal of the American Society for Horticultural Science* **120**, 808–813.
- Bukhov NG, Drozdova IS, Bondar VV. 1995. Light response curves of photosynthesis in leaves of sun-type and shade-type plants grown in blue or red light. *Journal of Photochemistry and Photobiology B: Biology* **30**, 39–41.
- Caird MA, Richards JH, Donovan LA. 2007. Nighttime stomatal conductance and transpiration in  $C_3$  and  $C_4$  plants. *Plant Physiology* **143**, 4–10.
- Christopher DA, Mullet JE. 1994. Separate photosensory pathways co-regulate blue-light/ultraviolet-A-activated *psbD*–*psbC* transcription and light-induced D2 and CP43 degradation in barley (*Hordeum vulgare*) chloroplasts. *Plant Physiology* **104**, 1119–1129.
- Evans JR, Poorter H. 2001. Photosynthetic acclimation of plants to growth irradiance: the relative importance of specific leaf area and nitrogen partitioning in maximizing carbon gain. *Plant, Cell and Environment* **24**, 755–767.
- Felippe GM, Dale JE, Marriott C. 1975. Effects of irradiance on uptake and assimilation of nitrate by young barley seedlings. *Annals of Botany* **39**, 43–55.
- Folta KM, Maruhnich SA. 2007. Green light: a signal to slow down or stop. *Journal of Experimental Botany* **58**, 3099–3111.

- Genty B, Briantais JM, Baker NR.** 1989. The relationship between the quantum yield of photosynthetic electron-transport and quenching of chlorophyll fluorescence. *Biochimica et Biophysica Acta* **990**, 87–92.
- Goins GD, Yorio NC, Sanwo MM, Brown CS.** 1997. Photomorphogenesis, photosynthesis, and seed yield of wheat plants grown under red light-emitting diodes (LEDs) with and without supplemental blue lighting. *Journal of Experimental Botany* **48**, 1407–1413.
- Harbinson J, Genty B, Baker NR.** 1990. The relationship between CO<sub>2</sub> assimilation and electron-transport in leaves. *Photosynthesis Research* **25**, 213–224.
- Hogewoning SW, Douwstra P, Trouwborst G, van Ieperen W, Harbinson J.** 2010. An artificial solar spectrum substantially alters plant development compared with usual climate room irradiance spectra. *Journal of Experimental Botany* **61**, 1267–1276.
- Hogewoning SW, Harbinson J.** 2007. Insights on the development, kinetics, and variation of photoinhibition using chlorophyll fluorescence imaging of a chilled, variegated leaf. *Journal of Experimental Botany* **58**, 453–463.
- Inada K.** 1976. Action spectra for photosynthesis in higher plants. *Plant and Cell Physiology* **17**, 355–365.
- Jarillo JA, Gabrys H, Capel J, Alonso JM, Ecker JR, Cashmore AR.** 2001. Phototropin-related NPL1 controls chloroplast relocation induced by blue light. *Nature* **410**, 952–954.
- Jarvis PG.** 1971. The estimation of resistance to carbon dioxide transfer. In: Sestak Z, Catsky J, Jarvis PG, eds. *Plant photosynthetic production: manual of methods*. The Hague: Junk, 566–631.
- Kingston-Smith AH, Harbinson J, Williams J, Foyer CH.** 1997. Effect of chilling on carbon assimilation, enzyme activation, and photosynthetic electron transport in the absence of photoinhibition in maize leaves. *Plant Physiology* **114**, 1039–1046.
- Lichtenthaler H, Buschmann C, Rahmsdorf U.** 1980. The importance of blue light for the development of sun-type chloroplasts. In: Senger H, ed. *The blue light syndrome*. Berlin: Springer-Verlag, 485–494.
- Massa GD, Kim HH, Wheeler RM, Mitchell CA.** 2008. Plant productivity in response to LED lighting. *Hortscience* **43**, 1951–1956.
- Matsuda R, Ohashi-Kaneko K, Fujiwara K, Goto E, Kurata K.** 2004. Photosynthetic characteristics of rice leaves grown under red light with or without supplemental blue light. *Plant and Cell Physiology* **45**, 1870–1874.
- Matsuda R, Ohashi-Kaneko K, Fujiwara K, Kurata K.** 2007. Analysis of the relationship between blue-light photon flux density and the photosynthetic properties of spinach (*Spinacia oleracea* L.) leaves with regard to the acclimation of photosynthesis to growth irradiance. *Soil Science and Plant Nutrition* **53**, 459–465.
- Matsuda R, Ohashi-Kaneko K, Fujiwara K, Kurata K.** 2008. Effects of blue light deficiency on acclimation of light energy partitioning in PSII and CO<sub>2</sub> assimilation capacity to high irradiance in spinach leaves. *Plant and Cell Physiology* **49**, 664–670.
- McCree KJ.** 1972. Action spectrum, absorptance and quantum yield of photosynthesis in crop plants. *Agricultural Meteorology* **9**, 191–216.
- Mochizuki T, Onda Y, Fujiwara E, Wada M, Toyoshima Y.** 2004. Two independent light signals cooperate in the activation of the plastid *psbD* blue light-responsive promoter in *Arabidopsis*. *FEBS Letters* **571**, 26–30.
- Nejad AR, van Meeteren U.** 2005. Stomatal response characteristics of *Tradescantia virginiana* grown at high relative air humidity. *Physiologia Plantarum* **125**, 324–332.
- Oguchi R, Hikosaka K, Hirose T.** 2003. Does the photosynthetic light-acclimation need change in leaf anatomy? *Plant, Cell and Environment* **26**, 505–512.
- Ohashi-Kaneko K, Matsuda R, Goto E, Fujiwara K, Kurata K.** 2006. Growth of rice plants under red light with or without supplemental blue light. *Soil Science and Plant Nutrition* **52**, 444–452.
- Paul MJ, Foyer CH.** 2001. Sink regulation of photosynthesis. *Journal of Experimental Botany* **52**, 1383–1400.
- Poorter H, Niinemets U, Poorter L, Wright IJ, Villar R.** 2009. Causes and consequences of variation in leaf mass per area (LMA): a meta-analysis. *New Phytologist* **182**, 565–588.
- Poorter H, Niinemets U, Walter A, Fiorani F, Schurr U.** 2010. A method to construct dose-response curves for a wide range of environmental factors and plant traits by means of a meta-analysis of phenotypic data. *Journal of Experimental Botany* **61**, 2043–2055.
- Sager JC, Smith WO, Edwards JL, Cyr KL.** 1988. Photosynthetic efficiency and phytochrome photoequilibria determination using spectral data. *Transactions of the ASAE* **31**, 1882–1889.
- Schmid R, Fromme R, Renger G.** 1990a. The photosynthetic apparatus of *Acetabularia mediterranea* grown under red or blue light. Biophysical quantification and characterization of photosystem II and its core components. *Photochemistry and Photobiology* **52**, 103–109.
- Schmid R, Wennecke R, Fleischhauer S.** 1990b. Quantitative correlation of peripheral and intrinsic core polypeptides of photosystem II with photosynthetic electron-transport activity of *Acetabularia mediterranea* in red and blue light. *Planta* **182**, 391–398.
- Sharkey TD, Raschke K.** 1981. Effect of light quality on stomatal opening in leaves of *Xanthium strumarium* L. *Plant Physiology* **68**, 1170–1174.
- Smith S, Weyers JDB, Berry WG.** 1989. Variation in stomatal characteristics over the lower surface of *Commelina communis* leaves. *Plant, Cell and Environment* **12**, 653–659.
- Snyder KA, Richards JH, Donovan LA.** 2003. Night-time conductance in C<sub>3</sub> and C<sub>4</sub> species: do plants lose water at night? *Journal of Experimental Botany* **54**, 861–865.
- Sood S, Tyagi AK, Tripathy BC.** 2004. Inhibition of photosystem I and photosystem II in wheat seedlings with their root-shoot transition zones exposed to red light. *Photosynthesis Research* **81**, 31–40.
- Stitt M.** 1991. Rising CO<sub>2</sub> levels and their potential significance for carbon flow in photosynthetic cells. *Plant, Cell and Environment* **14**, 741–762.
- Tennessen DJ, Singsaas EL, Sharkey TD.** 1994. Light-emitting diodes as a light-source for photosynthesis research. *Photosynthesis Research* **39**, 85–92.

- Thornley JHM.** 1976. *Mathematical models in plant physiology: a quantitative approach to problems in plant and crop physiology*. London: Academic Press.
- Trouwborst G, Oosterkamp J, Hogewoning SW, Harbinson J, van Ieperen W.** 2010. The responses of light interception, photosynthesis and fruit yield of cucumber to LED-lighting within the canopy. *Physiologia Plantarum* **138**, 289–300.
- von Caemmerer S, Farquhar GD.** 1981. Some relationships between the biochemistry of photosynthesis and the gas-exchange of leaves. *Planta* **153**, 376–387.
- Voskresenskaya NP, Drozdova IS, Krendeleva TE.** 1977. Effect of light quality on organization of photosynthetic electron-transport chain of pea-seedlings. *Plant Physiology* **59**, 151–154.
- Wellburn AR.** 1994. The spectral determination of chlorophyll a and chlorophyll b, as well as total carotenoids, using various solvents with spectrophotometers of different resolution. *Journal of Plant Physiology* **144**, 307–313.
- Wennicke H, Schmid R.** 1987. Control of the photosynthetic apparatus of *Acetabularia mediterranea* by blue light. Analysis by light-saturation curves. *Plant Physiology* **84**, 1252–1256.
- Whitelam G, Halliday K.** 2007. *Light and plant development*. Oxford: Blackwell Publishing.
- Yorio NC, Goins GD, Kagie HR, Wheeler RM, Sager JC.** 2001. Improving spinach, radish, and lettuce growth under red light-emitting diodes (LEDs) with blue light supplementation. *Hortscience* **36**, 380–383.
- Zheng S-J, Snoeren TAL, Hogewoning SW, van Loon JJA, Dicke M.** 2010. Disruption of plant carotenoid biosynthesis through virus-induced gene silencing affects oviposition behaviour of the butterfly *Pieris rapae*. *New Phytologist* **186**, 733–745.